



Theoretical Analysis of Dark Matter Distribution in Galactic Halos

¹Wanchai Leroux-Martin

¹Department of Electrical and Computer Engineering, Kavir Polytechnic University of Technology, Iran

¹wanchai.leroux.martin@kput-ir.net

Peer Review Information

Submission: 05 Jan 2021

Revision: 20 Feb 2021

Acceptance: 02 March 2021

Keywords

dark matter, galactic halos, NFW profile, Burkert profile, Einasto profile

Abstract

Dark matter halos dominate the gravitating mass of galaxies outside their luminous regions, yet the radial form of the halo density remains one of the central theoretical problems in galaxy dynamics. This methodology paper develops a controlled analytical comparison of three widely used spherical halo descriptions: the Navarro-Frenk-White (NFW), Burkert, and Einasto profiles. Each profile is normalized to an identical Milky-Way-scale virial mass, $M_{200} = 10^{12}$ solar masses, within $r_{200} = 210$ kpc so that differences in the inner halo arise from profile shape rather than total mass. The theoretical workflow evaluates density, enclosed mass, circular velocity, and logarithmic density slope over 0.1-210 kpc, with sensitivity tests on concentration, core radius, and Einasto shape index. Model-derived results show that the NFW and Einasto prescriptions remain centrally concentrated, whereas the Burkert model develops a nearly constant-density inner core. Despite marked differences at approximately 1 kpc, the profiles become more similar over intermediate galactic radii after common-mass normalization. The analysis demonstrates why high-resolution inner-galaxy kinematics are especially discriminating and provides a reproducible framework for testing halo structure without presenting illustrative calculations as observational measurements.

Introduction

Galaxy rotation curves and dynamical mass estimates require substantially more gravitating matter than can be supplied by stars and gas alone, motivating extended dark matter halos. Within the standard cold-dark-matter framework, collisionless structure formation naturally produces centrally concentrated halos. Early numerical studies converged on a nearly universal profile whose inner density increases approximately as r^{-1} and whose outer density approaches r^{-3} (Navarro, Frenk, & White, 1996, 1997). The resulting NFW profile became the baseline for theoretical halo modeling because it connects a characteristic radius and density to the virial mass and concentration of the halo.

The inner few kiloparsecs are, however, especially important. Observations of dwarf and low-surface-brightness systems have repeatedly

motivated cored alternatives in which the central density changes slowly with radius (Moore, 1994; Burkert, 1995). This difference between a central cusp and a core is not merely descriptive: it changes the enclosed mass, circular speed, gravitational potential, and predicted response of baryons. The core-cusp tension has therefore become a compact way to test the interaction between cosmological initial conditions, baryonic feedback, and possible non-standard dark-matter physics (de Blok, 2010; Bullock & Boylan-Kolchin, 2017).

A second development is that simulated halo profiles need not approach a fixed power-law slope at very small radius. High-resolution work found that the local logarithmic slope varies continuously with radius and can be represented well by Einasto-type forms (Navarro et al., 2004;

Merritt et al., 2006). This motivates a three-model comparison: NFW as the conventional cusped reference, Burkert as a phenomenological cored profile, and Einasto as a smoothly varying-slope profile. The aim of the present paper is not to fit a specific galaxy, but to formulate and demonstrate a theoretical method for comparing these halo families under identical global mass constraints.

Literature Review

The literature on halo structure developed through a sequence of increasingly precise theoretical and observational constraints. Moore (1994) highlighted that galaxy-halo observations could conflict with steep dissipationless dark-matter distributions, while Burkert (1995) proposed a finite-density empirical profile for dwarf-galaxy halos. These studies established the observational motivation for cored mass models. In parallel, Navarro, Frenk, and White (1996, 1997) demonstrated through cosmological N-body simulations that equilibrium halos over a wide mass range could be represented by a common scaled density law. Their formulation introduced the characteristic scale radius and concentration framework that remains central to halo phenomenology.

Improved numerical resolution subsequently showed that the inner slope is not necessarily a single asymptotic power law. Navarro et al. (2004) reported a progressive change of slope toward small radii and motivated curved density descriptions. Merritt et al. (2006) compared parametric forms with simulated halos and found strong performance for Einasto-like models. Large cosmological calculations such as the Aquarius Project extended this picture to the

hierarchy and internal substructure of Milky-Way-size halos, emphasizing that smooth radial structure coexists with surviving subhalos (Springel et al., 2008).

Theoretical halo profiles must ultimately be interpreted against galaxy-scale kinematics. de Blok (2010) reviewed the core-cusp problem and stressed that high-resolution velocity fields tend to favor shallower inner distributions in many gas-rich dwarf and low-surface-brightness galaxies. Oman et al. (2015) broadened the issue by showing that observed dwarf-galaxy rotation curves exhibit greater diversity at fixed characteristic velocity than simulated systems. Read et al. (2016) demonstrated with high-resolution dwarf simulations that repeated baryonic feedback can transform a cusp into a core under suitable star-formation histories, while Bullock and Boylan-Kolchin (2017) placed the core-cusp problem among the principal small-scale tests of the Lambda-CDM paradigm.

By 2021, attention had shifted increasingly toward quantifying modeling systematics and the amount of data required to distinguish profile shapes. Chang and Necib (2021), using mock observations and Jeans modeling, showed that robust inference of inner slopes in dwarf galaxies can require very large stellar samples. Their result is important for methodology: an apparent preference for a core or cusp can reflect sampling and dynamical assumptions as well as the true halo. Therefore, a clean theoretical comparison should first establish how much the candidate profiles differ under controlled normalization before observational uncertainties are introduced.

Table 1. Selected development of galactic-halo density modeling through 2021

Study	Year	Main contribution	Use in this paper
Moore	1994	Early evidence for a halo core/cusp discrepancy	Motivation for inner-profile testing
Burkert	1995	Finite-density cored halo model	Cored comparison profile
Navarro et al.	1996/1997	Universal cusped CDM profile	Baseline NFW model
Navarro et al.	2004	Continuously varying inner slope	Supports curved-slope modeling
Merritt et al.	2006	Einasto performs well for simulated halos	Einasto comparison profile
Springel et al.	2008	High-resolution Milky Way halo/subhalo structure	Context for CDM halo complexity

de Blok	2010	Review of the core-cusp problem	Observational interpretation
Chang & Necib	2021	Jeans-modeling systematics and sample-size limits	Caution in inference

Methodology

1. Research Design and Assumptions

The study uses a deterministic theoretical-computational design. No observed rotation-curve data are fitted and no new empirical measurement is claimed. The purpose is to isolate the structural consequences of alternative halo density laws under the same global halo mass. Spherical symmetry and dynamical equilibrium are assumed, and baryonic contributions from the stellar disk, bulge, and gas are excluded from the baseline calculation. These assumptions deliberately reduce the problem to the dark-matter component and make the interpretation of profile shape transparent.

A fiducial halo mass $M_{200} = 1.0 \times 10^{12}$ solar masses and virial radius $r_{200} = 210$ kpc are adopted. For NFW, the concentration is $c = 10$, giving $r_s = r_{200}/c = 21$ kpc. The Burkert core radius is $r_0 = 10$ kpc. The Einasto model uses $r_{-2} = 21$ kpc and shape parameter $\alpha = 0.18$. The characteristic density of each model is solved independently so that $M(<r_{200})$ is exactly 10^{12} solar masses. This common-mass normalization is essential: without it, differences in density could be caused by unequal halo mass rather than by the radial form of the model.

2. Density Models

The NFW profile is used as the standard cusped CDM reference (Navarro et al., 1996, 1997). With $x = r/r_s$, its density is

$$\rho_{\text{NFW}}(r) = \rho_s / [x(1+x)^2].$$

Its inner behavior is ρ proportional to r^{-1} and its outer behavior is ρ proportional to r^{-3} . Thus, the central density rises without forming a finite-density core.

The Burkert profile represents a cored halo (Burkert, 1995). With $x = r/r_0$,

$$\rho_B(r) = \rho_0 / [(1+x)(1+x^2)].$$

At r much smaller than r_0 , ρ_B approaches ρ_0 , producing an approximately constant-density center. At large radius the falloff tends toward r^{-3} , allowing a useful direct comparison with NFW while preserving different inner behavior.

The Einasto model is included because high-resolution simulations favor profiles whose

logarithmic slope changes smoothly with radius (Navarro et al., 2004; Merritt et al., 2006). Its adopted form is

$$\rho_E(r) = \rho_{-2} \exp\{-(2/\alpha)[(r/r_{-2})^\alpha - 1]\}.$$

Here r_{-2} is the radius where the local logarithmic slope equals -2 , and α controls the curvature of the profile. Unlike NFW, the model does not impose a fixed inner power-law asymptote over the resolved range.

3. Derived Dynamical Quantities

For each density law, enclosed mass is calculated from spherical integration,

$$M(<r) = 4\pi \int_0^r \rho(r') r'^2 dr'.$$

The corresponding circular velocity is then evaluated from $V_c(r) = [GM(<r)/r]^{1/2}$ using $G = 4.30091 \times 10^{-6}$ kpc (km/s)² solar-mass⁻¹. The local shape is summarized by $\gamma(r) = -d \ln \rho / d \ln r$; γ close to zero denotes a core, whereas γ near one denotes an NFW-like cusp.

4. Computational Procedure

The radial domain is sampled logarithmically from 0.1 to 210 kpc. A logarithmic grid is preferred because the diagnostic information is concentrated at small radii while the model must still extend to the virial boundary. For each grid point the density is evaluated directly, the enclosed mass is obtained by numerical quadrature, and the circular speed and logarithmic slope are derived. The maximum circular velocity V_{max} and its radius $r(V_{\text{max}})$ are identified numerically. Values at 1, 8, and 10 kpc are retained as compact inner- and intermediate-radius diagnostics.

The calculation proceeds in five reproducible steps: (1) set M_{200} and r_{200} ; (2) choose profile-shape parameters; (3) solve the characteristic density that enforces $M(<r_{200}) = M_{200}$; (4) evaluate $\rho(r)$, $M(<r)$, $V_c(r)$, and $\gamma(r)$ on the common radial grid; and (5) compare profiles using central density, inner mass, maximum circular velocity, and slope. This workflow can later be embedded in an observational likelihood by adding stellar and gas mass models and a covariance matrix for measured velocities.

Table 2. Fiducial theoretical halo parameters

Model	M200 (Msun)	Scale parameter	Shape parameter	Normalization
NFW	1.0×10^{12}	$r_s = 21$ kpc	$c = 10$	$\rho_{s,0} = 5.77 \times 10^6 \text{ Msun kpc}^{-3}$
Burkert	1.0×10^{12}	$r_0 = 10$ kpc	cored	$\rho_{0,0} = 34.50 \times 10^6 \text{ Msun kpc}^{-3}$
Einasto	1.0×10^{12}	$r_{-2} = 21$ kpc	$\alpha = 0.18$	$\rho_{-2,0} = 1.38 \times 10^6 \text{ Msun kpc}^{-3}$

5. Sensitivity and Interpretation Rules

A basic sensitivity analysis varies the NFW concentration over $c = 8$ -12, the Burkert core radius over $r_0 = 5$ -15 kpc, and the Einasto shape index over $\alpha = 0.15$ -0.25 while re-normalizing the total virial mass after each change. The primary interpretation rule is that a model is considered strongly core-like only when γ remains much smaller than one throughout the inner kiloparsec. Because baryons are omitted, the calculated V_c curves are halo-only quantities and should not be compared directly with a measured total rotation curve without adding the gravitational contributions of stars and gas. This separation prevents overinterpretation of the theoretical exercise.




Figure 1. Model-derived dark-matter density profiles after normalization to the same M200 = 10^{12} solar masses. The central divergence of NFW, finite Burkert core, and continuously curved Einasto form are most distinguishable at small radii.

Results

Density and Inner-Slope Comparison

The common-mass normalization produces a clear separation in the inner halo. At 1 kpc, the

NFW density is 110.4×10^6 solar masses kpc^{-3} , the Burkert density is 31.1×10^6 solar masses kpc^{-3} , and the Einasto density is 149.5×10^6 solar masses kpc^{-3} . The Burkert value is therefore only about 0.28 of the NFW value at this radius. This difference is a direct consequence of the core: the Burkert logarithmic slope at 1 kpc is $\gamma = 0.11$, compared with 1.09 for NFW and 1.16 for Einasto.

At 8 kpc, the three models are substantially closer: NFW gives 7.94×10^6 solar masses kpc^{-3} , Burkert 11.69×10^6 , and Einasto 8.10×10^6 solar masses kpc^{-3} . This convergence at intermediate radius explains why outer rotation data alone may have limited power to determine whether the central halo is cored or cusped. The strongest diagnostic leverage is obtained by resolving the inner region where the slope and enclosed mass respond most directly to the assumed density law, consistent with the methodological concerns emphasized in de Blok (2010) and Chang and Necib (2021).

Enclosed Mass and Circular Velocity

Within 10 kpc, the calculated dark-matter masses are 4.49×10^{10} solar masses for NFW, 5.51×10^{10} for Burkert, and 4.85×10^{10} for Einasto. Because the Burkert halo is normalized to the same virial mass but assigned a 10 kpc core scale, it redistributes mass differently across the inner-to-intermediate region and reaches a larger halo-only V_{max} in this particular fiducial parameterization. The computed maxima are approximately 172.5, 200.3, and 171.0 km/s, respectively. These numbers are not universal predictions; they are outputs of the stated parameter choices and are included to demonstrate the comparison procedure.

Table 3. Model-derived comparison for the fiducial halo

Model	$\rho(1 \text{ kpc})$ 10^6 Msun/kpc^3	$\gamma(1 \text{ kpc})$	$M(<10 \text{ kpc})$ 10^{10} Msun	V_{max} (km/s)	$r(V_{\text{max}})$ (kpc)	Interpretation
NFW	110.4	1.09	4.49	172.5	45.4	Cusped
Burkert	31.1	0.11	5.51	200.3	32.3	Cored

Einasto	149.5	1.16	4.85	171.0	45.9	Smoothly varying slope
---------	-------	------	------	-------	------	---------------------------

Conclusion and Discussion

This paper provides a compact theoretical methodology for analyzing dark-matter distribution in galactic halos using three widely adopted density laws. When NFW, Burkert, and Einasto halos are constrained to the same virial mass and radius, their most important differences occur in the central region rather than near the virial boundary. The NFW profile retains an r^{-1} -type cusp, the Burkert profile approaches a finite central density, and the Einasto profile changes slope continuously. The model-derived comparison therefore supports a central methodological conclusion: resolving the inner kiloparsec is much more informative for distinguishing profile families than relying only on intermediate- or outer-radius kinematics. The comparison also illustrates why parameter normalization must be handled carefully. A cored model can have a lower density at very small radius yet a different mass distribution at several kiloparsecs after the requirement of equal M_{200} is imposed. Circular-velocity differences are therefore controlled jointly by inner shape and global normalization. This is relevant to the observed diversity problem described by Oman et al. (2015): two galaxies with similar outer velocities need not have the same inner mass distribution, and a theoretical comparison must preserve that possibility rather than force all profiles to coincide at one inner point. The main limitation is deliberate simplicity. Real galaxies contain disks, bulges, gas, bars, triaxial structure, time-dependent feedback, and measurement uncertainties. Baryonic processes can alter the dark-matter distribution and may transform cusps into cores under suitable conditions, as shown in high-resolution dwarf simulations (Read et al., 2016). Conversely, dark-matter-only simulations retain rich substructure and centrally concentrated hosts (Springel et al., 2008). Thus, a discrepancy between a halo-only NFW curve and a measured rotation curve is not by itself evidence for new dark-matter physics. The luminous mass model and observational systematics must be included before such a conclusion is justified. Future work should extend this framework in three directions. First, the baryonic circular-speed terms should be added so that $V_{\text{total}}^2 = V_{\text{halo}}^2 + V_{\text{star}}^2 + V_{\text{gas}}^2 + V_{\text{bulge}}^2$ can be fitted to observed data. Second, Bayesian inference can be used to estimate posterior distributions for concentration, core radius, Einasto alpha, stellar mass-to-light ratio, and nuisance parameters.

Third, model comparison should use information criteria or Bayesian evidence rather than visual inspection alone. The 2021 literature on Jeans modeling further indicates that sampling density and kinematic precision must be treated as part of the inference problem rather than as secondary details (Chang & Necib, 2021). Under these extensions, the theoretical workflow developed here can serve as a transparent baseline for testing dark-matter halo structure in dwarf, spiral, and Milky-Way-like galaxies.

References

- Moore, B. (1994). Evidence against dissipation-less dark matter from observations of galaxy haloes. *Nature*, 370, 629-631. DOI: 10.1038/370629a0.
- Burkert, A. (1995). The structure of dark matter halos in dwarf galaxies. *The Astrophysical Journal Letters*, 447, L25-L28. DOI: 10.1086/309560.
- Navarro, J. F., Frenk, C. S., & White, S. D. M. (1996). The structure of cold dark matter halos. *The Astrophysical Journal*, 462, 563-575. DOI: 10.1086/177173.
- Navarro, J. F., Frenk, C. S., & White, S. D. M. (1997). A universal density profile from hierarchical clustering. *The Astrophysical Journal*, 490, 493-508. DOI: 10.1086/304888.
- Navarro, J. F., Hayashi, E., Power, C., Jenkins, A. R., Frenk, C. S., White, S. D. M., Springel, V., Stadel, J., & Quinn, T. R. (2004). The inner structure of LambdaCDM haloes - III. Universality and asymptotic slopes. *Monthly Notices of the Royal Astronomical Society*, 349, 1039-1051. DOI: 10.1111/j.1365-2966.2004.07586.x.
- Merritt, D., Graham, A. W., Moore, B., Diemand, J., & Terzic, B. (2006). Empirical models for dark matter halos. I. Nonparametric construction of density profiles and comparison with parametric models. *The Astronomical Journal*, 132, 2685-2700. DOI: 10.1086/508988.
- Springel, V., Wang, J., Vogelsberger, M., Ludlow, A., Jenkins, A., Helmi, A., Navarro, J. F., Frenk, C. S., & White, S. D. M. (2008). The Aquarius Project: the subhaloes of galactic haloes. *Monthly Notices of the Royal Astronomical Society*, 391, 1685-1711. DOI: 10.1111/j.1365-2966.2008.14066.x.

de Blok, W. J. G. (2010). The core-cusp problem. *Advances in Astronomy*, 2010, 789293. DOI: 10.1155/2010/789293.

Oman, K. A., Navarro, J. F., Fattahi, A., Frenk, C. S., Sawala, T., White, S. D. M., et al. (2015). The unexpected diversity of dwarf galaxy rotation curves. *Monthly Notices of the Royal Astronomical Society*, 452, 3650-3665. DOI: 10.1093/mnras/stv1504.

Read, J. I., Agertz, O., & Collins, M. L. M. (2016). Dark matter cores all the way down. *Monthly*

Notices of the Royal Astronomical Society, 459, 2573-2590. DOI: 10.1093/mnras/stw713.

Bullock, J. S., & Boylan-Kolchin, M. (2017). Small-scale challenges to the Λ CDM paradigm. *Annual Review of Astronomy and Astrophysics*, 55, 343-387. DOI: 10.1146/annurev-astro-091916-055313.

Chang, L. J., & Necib, L. (2021). Dark matter density profiles in dwarf galaxies: linking Jeans modelling systematics and observation. *Monthly Notices of the R*